

The Topology of Chaos



Robert Gilmore

Physics Department
Drexel University
Philadelphia, PA 19104
robert.gilmore@drexel.edu

Colloquium, Physics Department
University of Florida, Gainesville, FL

October 6, 2008

The Topology of Chaos



Robert Gilmore

Physics Department

Drexel University

Philadelphia, PA 19104

robert.gilmore@drexel.edu

Colloquium, Physics Department
University of Florida, Gainesville, FL
October 23, 2008

Outline


- 1 Overview
- 2 Experimental Challenge
- 3 Topology of Orbits
- 4 Topological Analysis Program
- 5 Basis Sets of Orbits
- 6 Bounding Tori
- 7 Covers and Images
- 8 Quantizing Chaos
- 9 Representation Theory of Strange Attractors
- 10 Summary

J. R. Tredicce

Can you explain my data?

I dare you to explain my data!

Where is Tredicce coming from?




Feigenbaum:

$$\alpha = 4.66920 16091 \dots$$

$$\delta = -2.50290 78750 \dots$$

Laser with Modulated Losses Experimental Arrangement



Original Objectives

Construct a simple, algorithmic procedure for:

- Classifying strange attractors
- Extracting classification information

from experimental signals.

Result

There is now a classification theory.

- ① It is topological
- ② It has a hierarchy of 4 levels
- ③ Each is discrete
- ④ There is rigidity and degrees of freedom
- ⑤ It is applicable to R^3 only — for now

The 4 Levels of Structure

- **Basis Sets of Orbits**
- **Branched Manifolds**
- **Bounding Tori**
- **Extrinsic Embeddings**

Organization

LINKS OF PERIODIC ORBITS

organize

BOUNDING TORI

organize

BRANCHED MANIFOLDS

organize


LINKS OF PERIODIC ORBITS

Experimental Schematic

The Topology
of Chaos

Robert
Gilmore

Laser Experimental Arrangement




Experimental Motivation

The Topology
of Chaos

Robert
Gilmore

Oscilloscope Traces




Results, Single Experiment

The Topology
of Chaos

Robert
Gilmore

Bifurcation Schematics




Some Attractors

The Topology
of Chaos

Robert
Gilmore

Coexisting Basins of Attraction




Many Experiments



The Topology
of Chaos

Robert
Gilmore

Bifurcation Perestroikas




Experimental Data: LSA




Lefranc - Cargese


Experimental Data: LSA



Stretching & Squeezing in a Torus



Rotating the Poincaré Section around the axis of the torus



Rotating the Poincaré Section around the axis of the torus





Figure 2. Left: Intersections of a chaotic attractor with a series of section planes are computed. Right: Their evolution from plane to plane shows the interplay of the stretching and squeezing mechanisms.

Another Visualization

The Topology
of Chaos






Robert
Gilmore

Cutting Open a Torus




Satisfying Boundary Conditions

Global Torsion




(a)



(b)

A Chemical Experiment

The Belousov-Zhabotinskii Reaction



Chaos

Motion that is


- **Deterministic:** $\frac{dx}{dt} = f(x)$
- **Recurrent**
- **Non Periodic**
- **Sensitive to Initial Conditions**

Strange Attractor

The Ω limit set of the flow. There are unstable periodic orbits “in” the strange attractor. They are

- “Abundant”
- Outline the Strange Attractor
- Are the Skeleton of the Strange Attractor

UPOs Outline Strange attractors



UPOs Outline Strange attractors




Figure 5. Left: a chaotic attractor reconstructed from a time series from a chaotic laser ; Right : Superposition of 12 periodic orbits of periods from 1 to 10.

Organization of UPOs in R^3 : Gauss Linking Number

$$LN(A, B) = \frac{1}{4\pi} \oint \oint \frac{(\mathbf{r}_A - \mathbf{r}_B) \cdot d\mathbf{r}_A \times d\mathbf{r}_B}{|\mathbf{r}_A - \mathbf{r}_B|^3}$$

Interpretations of $LN \simeq$ # Mathematicians in World

Linking Number of Two UPOs





Figure 6. Left: two periodic orbits of periods 1 and 4 embedded in a strange attractor; Right: a link of two knots that is equivalent to the pair of periodic orbits up to continuous deformations without crossings.

One Stretch-&-Squeeze Mechanism




Motion of Blobs in Phase Space

The Topology
of Chaos

Robert
Gilmore


Stretching — Squeezing



Birman - Williams Projection

Identify x and y if

$$\lim_{t \rightarrow \infty} |x(t) - y(t)| \rightarrow 0$$



Birman - Williams Theorem

If:

Then:

Birman - Williams Theorem

If: **Certain Assumptions**

Then:

Birman - Williams Theorem

If: **Certain Assumptions**

Then: **Specific Conclusions**

Assumptions, B-W Theorem

A flow $\Phi_t(x)$

- on R^n is dissipative, $n = 3$, so that $\lambda_1 > 0, \lambda_2 = 0, \lambda_3 < 0$.
- Generates a hyperbolic strange attractor SA

IMPORTANT: The underlined assumptions can be relaxed.

Conclusions, B-W Theorem

- The projection maps the strange attractor \mathcal{SA} onto a 2-dimensional branched manifold \mathcal{BM} and the flow $\Phi_t(x)$ on \mathcal{SA} to a semiflow $\bar{\Phi}(x)_t$ on \mathcal{BM} .
- UPOs of $\Phi_t(x)$ on \mathcal{SA} are in 1-1 correspondence with UPOs of $\bar{\Phi}(x)_t$ on \mathcal{BM} . Moreover, every link of UPOs of $(\Phi_t(x), \mathcal{SA})$ is isotopic to the correspond link of UPOs of $(\bar{\Phi}(x)_t, \mathcal{BM})$.

Remark: "One of the few theorems useful to experimentalists."


A Very Common Mechanism

The Topology
of Chaos


Robert
Gilmore

Rössler:

Attractor



Branched Manifold




A Mechanism with Symmetry

The Topology
of Chaos


Robert
Gilmore

Lorenz:

Attractor



Branched Manifold




Examples of Branched Manifolds

The Topology
of Chaos


Robert
Gilmore

Inequivalent Branched Manifolds


(a)




(b)



(c)



(d)





Aufbau Princip for Branched Manifolds

The Topology
of Chaos

Robert
Gilmore

Any branched manifold can be built up from stretching and squeezing units



subject to the conditions:

- Outputs to Inputs
- No Free Ends

Rössler System


(a) Rössler Equations

$$\frac{dx}{dt} = -y - z$$

$$\frac{dy}{dt} = x + ay$$

$$\frac{dz}{dt} = b + z(x - c)$$

(b)



(c)




(f)

$$\begin{pmatrix} -1 & 0 \\ 0 & 0 \end{pmatrix}$$

$$\begin{pmatrix} 0 & +1 \end{pmatrix}$$

(e)



(f)




Lorenz System

(a) Lorenz Equations

$$\frac{dx}{dt} = -\sigma x + \sigma y$$

$$\frac{dy}{dt} = \rho x - y - xz$$

$$\frac{dz}{dt} = -bz + xy$$




(f)


$$\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

$$\begin{pmatrix} + & - \\ - & + \end{pmatrix}$$

(e)



(d)



Poincaré Smiles at U s in R^3

- Determine organization of UPOs \Rightarrow
- Determine branched manifold \Rightarrow
- Determine equivalence class of \mathcal{SA}

Topological Analysis Program

Locate Periodic Orbits

Create an Embedding

Determine Topological Invariants (LN)


Identify a Branched Manifold

Verify the Branched Manifold

Model the Dynamics

Validate the Model

Method of Close Returns



Embeddings

Many Methods: Time Delay, Differential, Hilbert Transforms, SVD, Mixtures, ...

Tests for Embeddings: Geometric, Dynamic, Topological[†]

None Good

We Demand a 3 Dimensional Embedding

An Embedding and Periodic Orbits




Figure 5. Left: a chaotic attractor reconstructed from a time series from a chaotic laser ; Right : Superposition of 12 periodic orbits of periods from 1 to 10.

Determine Topological Invariants

The Topology
of Chaos

Robert
Gilmore

Linking Number of Orbit Pairs




Figure 6. Left: two periodic orbits of periods 1 and 4 embedded in a strange attractor; Right: a link of two knots that is equivalent to the pair of periodic orbits up to continuous deformations without crossings.

Lefranc - Cargese

Determine Topological Invariants

The Topology
of Chaos

Robert
Gilmore

Compute Table of Expt'l LN

Table 7.2 Linking numbers for all the surrogate periodic orbits, to period 8, extracted from Belousov-Zhabotinskii data^a

Orbit	Symbolics	1	2	3	4	5	6	7	8a	8b
1	1	0	1	1	2	2	2	3	4	3
2	01	1	1	2	3	4	4	5	6	6
3	011	1	2	2	4	5	6	7	8	8
4	0111	2	3	4	5	8	8	11	13	12
5	01 011	2	4	5	8	8	10	13	16	15
6	011 0M1	2	4	6	8	10	9	14	16	16
7	01 01 011	3	5	7	11	13	14	16	21	21
8a	01 01 0111	4	6	8	13	16	16	21	23	24
8b	01 011 011	3	6	8	12	15	16	21	24	21

^aAll indices are negative.

Determine Topological Invariants

Compare w. LN From Various BM

Table 2.1 Linking numbers for orbits to period five in Smale horseshoe dynamics.

	1^s	1^f	2_1	3^f	3^s	4_1	4_2^f	4_2^s	5_2^f	5_2^s	5_2^f	5_2^s	5_1^f	5_1^s
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	0	1	1	1	2	1	1	1	1	2	2	2	2
01	0	1	1	2	2	3	2	2	2	2	3	3	4	4
001	0	1	2	2	3	4	3	3	3	3	4	4	5	5
011	0	1	2	3	2	4	3	3	3	3	5	5	5	5
0111	0	2	3	4	4	5	4	4	4	4	7	7	8	8
0001	0	1	2	3	3	4	3	4	4	4	5	5	5	5
0011	0	1	2	3	3	4	4	3	4	4	5	5	5	5
00001	0	1	2	3	3	4	4	4	4	5	5	5	5	5
00011	0	1	2	3	3	4	4	4	5	4	5	5	5	5
00111	0	2	3	4	5	7	5	5	5	5	6	7	8	9
00101	0	2	3	4	5	7	5	5	5	5	7	6	8	9
01101	0	2	4	5	5	8	5	5	5	5	8	8	8	10
01111	0	2	4	5	5	8	5	5	5	5	9	9	10	8

Guess Branched Manifold




Figure 7. “Combing” the intertwined periodic orbits (left) reveals their systematic organization (right) created by the stretching and squeezing mechanisms.

Lefranc - Cargese


Identification & 'Confirmation'

- \mathcal{BM} Identified by LN of small number of orbits
- Table of LN GROSSLY overdetermined
- Predict LN of additional orbits
- Rejection criterion

Determine Topological Invariants

What Do We Learn?

- \mathcal{BM} Depends on Embedding
- Some things depend on embedding, some don't
- Depends on Embedding: Global Torsion, Parity, ..
- Independent of Embedding: Mechanism



Perestroikas of Strange Attractors

The Topology
of Chaos

Robert
Gilmore

Evolution Under Parameter Change





Figure 11. Various templates observed in two laser experiments. Top left: schematic representation of the parameter space of forced nonlinear oscillators showing resonance tongues. Right: templates observed in the fiber laser experiment: global torsion increases systematically from one tongue to the next [40]. Bottom left: templates observed in the YAG laser experiment (only the branches are shown): there is a variation in the topological organization across one chaotic tongue [39, 41].

Perestroikas of Strange Attractors

The Topology
of Chaos

Robert
Gilmore

Evolution Under Parameter Change



Lefranc - Cargese

Analysis of Nonstationary Data





Figure 16. Top left: time series from an optical parametric oscillator showing a burst of irregular behavior. Bottom left: segment of the time series containing a periodic orbit of period 9. Right: embedding of the periodic orbit in a reconstructed phase space and representation of the braid realized by the orbit. The braid entropy is $h_T = 0.377$, showing that the underlying dynamics is chaotic. Reprinted from [61].

Model the Dynamics

A hodgepodge of methods exist: # Methods \simeq # Physicists

Validate the Model

Needed: Nonlinear analog of χ^2 test. OPPORTUNITY:
Tests that depend on entrainment/synchronization.



Compare with Original Objectives

Construct a simple, algorithmic procedure for:

- Classifying strange attractors
- Extracting classification information


from experimental signals.

Orbits Can be “Pruned”


The Topology
of Chaos

Robert
Gilmore

There Are Some Missing Orbits




Lorenz



Shimizu-Morioka

Orbit Forcing



$A \Rightarrow B$

$B \Rightarrow C$


$A \Rightarrow C$

An Ongoing Problem


The Topology
of Chaos

Robert
Gilmore

Forcing Diagram - Horseshoe



U - SEQUENCE ORDER



Status of Problem

- Horseshoe organization - active
- More folding - barely begun
- Circle forcing - even less known
- Higher genus - new ideas required

Constraints on Branched Manifolds


“Inflate” a strange attractor

Union of ϵ ball around each point

Boundary is surface of bounded 3D manifold

Torus that bounds strange attractor

Torus, Longitudes, Meridians



Surface Singularities

Flow field: three eigenvalues: +, 0, -

Vector field “perpendicular” to surface

Eigenvalues on surface at fixed point: +, -

All singularities are regular saddles


$$\sum_{s.p.} (-1)^{\text{index}} = \chi(S) = 2 - 2g$$

fixed points on surface = index = $2g - 2$


Flow Near a Singularity




(a)



(b)




(c)




(d)


Torus Bounding Lorenz-like Flows




Twisting the Lorenz Attractor




(a)



(c)



(b)




(d)

Constraints Provided by Bounding Tori

The Topology
of Chaos


Robert
Gilmore

**Two possible branched manifolds
in the torus with $g=4$.**



(a)

(b)



(c)

Bounding Tori contain all known Strange Attractors

Tab.1. All known strange attractors of dimension $d_L < 3$ are bounded by one of the standard dressed tori.

Strange Attractor	Dressed Torus	Period $g - 1$ Orbit
Rossler, Duffing, Burke and Shaw	A_1	1
Various Lasers, Gateau Roule	A_1	1
Neuron with Subthreshold Oscillations	A_1	1
Shaw-van der Pol	$A_1 \cup A_1^{(1)}$	$1 \cup 1$
Lorenz, Shimizu-Morioka, Rikitake	A_2	$(12)^2$
Multispiral attractors	A_n	$(12^{n-1})^2$
C_n Covers of Rossler	C_n	1^n
C_2 Cover of Lorenz ^(a)	C_4	1^4
C_2 Cover of Lorenz ^(b)	A_8	$(122)^2$
C_n Cover of Lorenz ^(a)	C_{2n}	1^{2n}
C_n Cover of Lorenz ^(b)	P_{n+1}	$(1n)^n$
$2 \rightarrow 1$ Image of Fig. 8 Branched Manifold	A_8	$(122)^2$
Fig. 8 Branched Manifold	P_8	$(14)^4$

^(a) Rotation axis through origin.
^(b) Rotation axis through one focus.

Labeling Bounding Tori

Poincaré section is disjoint union of $g-1$ disks

Transition matrix sum of two $g-1 \times g-1$ matrices

One is cyclic $g-1 \times g-1$ matrix

Other represents union of cycles

Labeling via (permutation) group theory

Some Bounding Tori

The Topology
of Chaos




Robert
Gilmore

Bounding Tori of Low Genus

TABLE I: Enumeration of canonical forms up to genus 9

g	m	(p_1, p_2, \dots, p_m)	$n_1 n_2 \dots n_{g-1}$
1	1	(0)	1
3	2	(2)	11
4	3	(3)	111
5	4	(4)	1111
5	3	(2,2)	1212
6	5	(5)	11111
6	4	(3,2)	12112
7	6	(6)	111111
7	5	(4,2)	112121
7	5	(3,3)	112112
7	4	(2,2,2)	122122
7	4	(2,2,2)	131313
8	7	(7)	1111111
8	6	(5,2)	1211112
8	6	(4,3)	1211121
8	5	(3,2,2)	1212212
8	5	(3,2,2)	1221221
8	5	(3,2,2)	1313131
9	8	(8)	11111111
9	7	(6,2)	11111212
9	7	(5,3)	11112112
9	7	(4,4)	11121112
9	6	(4,2,2)	11122122
9	6	(4,2,2)	11131313
9	6	(4,2,2)	11212212
9	6	(4,2,2)	12121212
9	6	(3,3,2)	11212122
9	6	(3,3,2)	11221122
9	6	(3,3,2)	11221212
9	6	(3,3,2)	11313133
9	5	(2,2,2,2)	12221222
9	5	(2,2,2,2)	12313132
9	5	(2,2,2,2)	14141414

Some Genus-9 Bounding Tori




Aufbau Princip for Bounding Tori

The Topology
of Chaos

Robert
Gilmore

Any bounding torus can be built up from equal numbers of stretching and squeezing units




- **Outputs to Inputs**
- **No Free Ends**
- **Colorless**

Aufbau Princip for Bounding Tori

The Topology
of Chaos

Robert
Gilmore

Application: Lorenz Dynamics, $g=3$



Construction of Poincaré Section

P. S. = Union 

Components = $g-1$

The Growth is Exponential


TABLE I: Number of canonical bounding tori as a function of genus, g .

g	$N(g)$	g	$N(g)$	g	$N(g)$
3	1	9	15	15	2211
4	1	10	28	16	5549
5	2	11	67	17	14290
6	2	12	145	18	36824
7	5	13	368	19	96347
8	6	14	870	20	252927

The Growth is Exponential The Entropy is $\log 3$

Bounding Torus Entropy

$\text{Log}[N(g)]/(g-1)$




Extrinsic Embedding of Bounding Tori

The Topology
of Chaos

Robert
Gilmore

Extrinsic Embedding of Intrinsic Tori



Partial classification by links of homotopy group generators.
Nightmare Numbers are Expected.



Modding Out a Rotation Symmetry

The Topology
of Chaos

Robert
Gilmore

Modding Out a Rotation Symmetry





$$\begin{pmatrix} X \\ Y \\ Z \end{pmatrix} \rightarrow \begin{pmatrix} u \\ v \\ w \end{pmatrix} = \begin{pmatrix} \operatorname{Re} (X + iY)^2 \\ \operatorname{Im} (X + iY)^2 \\ Z \end{pmatrix}$$



Lorenz Attractor and Its Image

The Topology
of Chaos

Robert
Gilmore





Lifting an Attractor: Cover-Image Relations

The Topology
of Chaos


Robert
Gilmore

Creating a Cover with Symmetry


$$\begin{pmatrix} X \\ Y \\ Z \end{pmatrix} \leftarrow \begin{pmatrix} u \\ v \\ w \end{pmatrix} = \begin{pmatrix} \operatorname{Re} (X + iY)^2 \\ \operatorname{Im} (X + iY)^2 \\ Z \end{pmatrix}$$




Cover-Image Branched Manifolds



Two Two-fold Lifts Different Symmetry




**Rotation
Symmetry**




**Inversion
Symmetry**

Topological Index: Choose Group Choose Rotation Axis (Singular Set)



Different Rotation Axes Produce Different (Nonisotopic) Lifts



Nonisotopic Locally Diffeomorphic Lifts

The Topology
of Chaos


Robert
Gilmore



(a) $\mu = 0.0$



(c) $\mu = -2.083$




(e) $\mu = -4.166$



(b) $\mu = -0.84548$



(d) $\mu = -3.14674$





Indices $(0,1)$ and $(1,1)$


The Topology
of Chaos

Robert
Gilmore

Two Two-fold Covers Same Symmetry









Three-fold, Four-fold Covers



Two Inequivalent Lifts with V_4 Symmetry

The Topology
of Chaos

Robert
Gilmore



Algorithm

- Construct Invariant Polynomials, Syzygies, Radicals
- Construct Singular Sets
- Determine Topological Indices
- Construct Spectrum of Structurally Stable Covers
- Structurally Unstable Covers Interpolate


Symmetries Due to Symmetry

- Schur's Lemmas & Equivariant Dynamics
- Cauchy Riemann Symmetries
- Clebsch-Gordon Symmetries
- Continuations
 - Analytic Continuation
 - Topological Continuation
 - Group Continuation


Covers of a Trefoil Torus

The Topology
of Chaos


Robert
Gilmore



Granny Knot



Square Knot




Trefoil Knot

You Can Cover a Cover = Lift a Lift


The Topology
of Chaos

Robert
Gilmore


Covers of Covers of Covers





Rossler



Lorenz



Every Knot Lives Here



Local Stuff

Groups:

Local Isomorphisms


Cartan's Theorem

Dynamical Systems:

Local Diffeomorphisms

??? Anything Useful ???

Cartan's Theorem for Lie Groups




Universal Image Dynamical System

The Topology
of Chaos

Robert
Gilmore

Locally Diffeomorphic Covers of \underline{D}



\underline{D} : Universal Image Dynamical System

Local Isomorphisms & Diffeomorphisms

Local Isomorphisms & Diffeomorphisms

Lie Groups

Local Isomorphisms & Diffeomorphisms


Lie Groups

Local Isomorphisms

Local Isomorphisms & Diffeomorphisms

Lie Groups

Local Isomorphisms




Local Isomorphisms & Diffeomorphisms

Lie Groups

Dynamical Systems

Local Isomorphisms




Local Isomorphisms & Diffeomorphisms

Lie Groups

Dynamical Systems

Local Isomorphisms

Local Diffeos





Local Isomorphisms & Diffeomorphisms

Lie Groups

Dynamical Systems

Local Isomorphisms

Local Diffeos



Rotating the Attractor

$$\frac{d}{dt} \begin{bmatrix} X \\ Y \end{bmatrix} = \begin{bmatrix} F_1(X, Y) \\ F_2(X, Y) \end{bmatrix} + \begin{bmatrix} a_1 \sin(\omega_d t + \phi_1) \\ a_2 \sin(\omega_d t + \phi_2) \end{bmatrix}$$

$$\begin{bmatrix} u(t) \\ v(t) \end{bmatrix} = \begin{bmatrix} \cos \Omega t & -\sin \Omega t \\ \sin \Omega t & \cos \Omega t \end{bmatrix} \begin{bmatrix} X(t) \\ Y(t) \end{bmatrix}$$

$$\frac{d}{dt} \begin{bmatrix} u \\ v \end{bmatrix} = R\mathbf{F}(R^{-1}\mathbf{u}) + R\mathbf{t} + \Omega \begin{bmatrix} -v \\ +u \end{bmatrix}$$

$$\Omega = n \omega_d$$

$$q \Omega = p \omega_d$$

Global Diffeomorphisms

Local Diffeomorphisms
(p -fold covers)


Two Phase Spaces: R^3 and $D^2 \times S^1$

The Topology
of Chaos


Robert
Gilmore

Rosler Attractor: Two Representations

R^3





$D^2 \times S^1$



Rossler Attractor:

Two More Representations with $n = \pm 1$





Subharmonic, Locally Diffeomorphic Attractors

The Topology
of Chaos

Robert
Gilmore

Rossler Attractor:

Two Two-Fold Covers with $p/q = \pm 1/2$





Subharmonic, Locally Diffeomorphic Attractors

The Topology
of Chaos

Robert
Gilmore

Rosler Attractor:

Two Three-Fold Covers with $p/q = -2/3, -1/3$





Subharmonic, Locally Diffeomorphic Attractors

The Topology
of Chaos

Robert
Gilmore

Rossler Attractor:

And Even More Covers (with $p/q = +1/3, +2/3$)



Angular Momentum and Energy

$$L(0) = \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int_0^\tau X dY - Y dX \qquad K(0) = \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int_0^\tau \frac{1}{2} (\dot{X}^2 + \dot{Y}^2) dt$$

$$L(\Omega) = \langle uv - vu \rangle \qquad K(\Omega) = \left\langle \frac{1}{2} (\dot{u}^2 + \dot{v}^2) \right\rangle$$

$$= L(0) + \Omega \langle R^2 \rangle \qquad = K(0) + \Omega L(0) + \frac{1}{2} \Omega^2 \langle R^2 \rangle$$

$$\langle R^2 \rangle = \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int_0^\tau (X^2 + Y^2) dt = \lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int_0^\tau (u^2 + v^2) dt$$

New Measures, Diffeomorphic Attractors


The Topology
of Chaos

Robert
Gilmore


Energy and Angular Momentum

Diffeomorphic, Quantum Number n

Torsion Integral



Energy Integral




New Measures, Subharmonic Covering Attractors

The Topology
of Chaos


Robert
Gilmore

Energy and Angular Momentum Subharmonics, Quantum Numbers p/q

Torsion Integral



Energy Integral



Embeddings

An embedding creates a diffeomorphism between an ('invisible') dynamics in someone's laboratory and a ('visible') attractor in somebody's computer.

Embeddings provide a *representation* of an attractor.

Equivalence is by Isotopy.

Irreducible is by Dimension

Inequivalent Irreducible Representations

Irreducible Representations of 3-dimensional Genus-one attractors are distinguished by three topological labels:

Parity	P
Global Torsion	N
Knot Type	KT

$$\Gamma^{P,N,KT}(\mathcal{SA})$$

Mechanism (stretch & fold, stretch & roll) is an invariant of embedding. It is independent of the representation labels.

Equivalent Reducible Representations

Topological indices (P,N,KT) are obstructions to isotopy for embeddings of minimum dimension (irreducible representations).

Are these obstructions removed by injections into higher dimensions (reducible representations)?

Systematically?

Equivalences by Injection

Obstructions to Isotopy

$$\begin{array}{ccc} R^3 & \longrightarrow & R^4 & \longrightarrow & R^5 \\ \text{Global Torsion} & & \text{Global Torsion} & & \\ \text{Parity} & & & & \\ \text{Knot Type} & & & & \end{array}$$

There is one *Universal* reducible representation in R^N , $N \geq 5$.
In R^N the only topological invariant is *mechanism*.

Summary

**1 Question Answered \Rightarrow
2 Questions Raised**

We must be on the right track !

Original Objectives Achieved

There is now a simple, algorithmic procedure for:

- Classifying strange attractors
- Extracting classification information

from experimental signals.

Result

There is now a classification theory for low-dimensional strange attractors.

- ① It is topological
- ② It has a hierarchy of 4 levels
- ③ Each is discrete
- ④ There is rigidity and degrees of freedom
- ⑤ It is applicable to R^3 only — for now

The Classification Theory has 4 Levels of Structure

The Classification Theory has 4 Levels of Structure

- 1 Basis Sets of Orbits

The Classification Theory has 4 Levels of Structure

- 1 Basis Sets of Orbits
- 2 Branched Manifolds

The Classification Theory has 4 Levels of Structure

- 1 Basis Sets of Orbits
- 2 Branched Manifolds
- 3 Bounding Tori



The Classification Theory has 4 Levels of Structure

- 1 Basis Sets of Orbits
- 2 Branched Manifolds
- 3 Bounding Tori
- 4 Extrinsic Embeddings

Four Levels of Structure

The Topology
of Chaos

Robert
Gilmore



Poetic Organization

LINKS OF PERIODIC ORBITS

organize

BOUNDING TORI

organize

BRANCHED MANIFOLDS

organize

LINKS OF PERIODIC ORBITS

Some Unexpected Results

- Perestroikas of orbits constrained by branched manifolds
- Routes to Chaos = Paths through orbit forcing diagram
- Perestroikas of branched manifolds constrained by bounding tori
- Global Poincaré section = union of $g - 1$ disks
- Systematic methods for cover - image relations
- Existence of topological indices (cover/image)
- Universal image dynamical systems
- NLD version of Cartan's Theorem for Lie Groups
- Topological Continuation – Group Continuation
- Cauchy-Riemann symmetries
- Quantizing Chaos
- Representation labels for inequivalent embeddings
- Representation Theory for Strange Attractors

We hope to find:

- Robust topological invariants for R^N , $N > 3$
- A Birman-Williams type theorem for higher dimensions
- An algorithm for irreducible embeddings
- Embeddings: better methods and tests
- Analog of χ^2 test for NLD
- Better forcing results: Smale horseshoe, $D^2 \rightarrow D^2$,
 $n \times D^2 \rightarrow n \times D^2$ (e.g., Lorenz), $D^N \rightarrow D^N$, $N > 2$
- Representation theory: complete
- Singularity Theory: Branched manifolds, splitting points
(0 dim.), branch lines (1 dim).
- Singularities as obstructions to isotopy