

Name: _____

Multiple Choice

Choose the better choice of all choices given.

1. 2 points How many electrons can fit in one electron orbital shell?
 - A. Only one
 - B. Only two
 - C. Only three
 - D. Unlimited.

2. 2 points Which of the following isn't a truth about quantum mechanics?
 - A. Energy is quantized
 - B. Particles have wavelike properties.
 - C. A cat can be dead and alive at the same time.
 - D. A particle can be in a combination of two different states.

3. 2 points Given what we've learned about relativity, which of the following is better to say than "matter can neither be created nor destroyed":
 - A. Matter can be converted to waves.
 - B. Matter can be converted to particles.
 - C. Matter exists as a particle and wave at the same time.
 - D. Mass and energy can be seen as two names for the same underlying, conserved physical quantity.

4. 2 points Which of the following problem in physics wasn't fixed by quantum mechanics?
 - A. The ultraviolet catastrophe of blackbody radiation
 - B. The photoelectric mystery
 - C. The helium spectrum mystery
 - D. The contradiction between the universal speed of light and Galilean transforms.

5. 2 points Planck's quantization of energy was a desperate attempt to resolve the
 - A. twin paradox.
 - B. ultraviolet catastrophe.
 - C. photoelectric mystery.
 - D. spectrum of hydrogen mystery.

Please go on to the next page...

-
6. 2 points We don't experience energy quantization in the everyday world because at our size
- A. the spaces between energy levels is too small to see with our senses.
 - B. quantum mechanics isn't true on the macroscopic scale.
 - C. energy becomes continuous past $n = 493, 523, 235, 674, 453, 213, 956$.
7. 2 points If I know the velocity of a subatomic particle precisely, then
- A. I know nothing about the particle's position.
 - B. I know a very limited amount about the particle's position.
 - C. The particle must be at rest.
 - D. The particle can't be at rest.

Problems

8. 5 points An HCL molecule vibrates with a frequency of 8.1×10^{13} Hz. What is the smallest possible change of energy that this molecule could experience?
9. 5 points How much energy is needed to ionize a hydrogen atom in the $n = 2$ state? What frequency photon would that require?

10. 5 points What percentage of mass does a hydrogen atom lose when it transitions from the $n = 3$ to the $n = 1$ state (in other words, $\Delta m/m_0$)? The rest mass of hydrogen is $m_0 c^2 = 939 \times 10^6$ eV.
11. Scientist A measures a quantum system by injecting an electron into a potential well, measuring its position with a laser beam (by finding where the photons reflect using Compton's equation, for example). After each measurement of its position, Scientist A flushes the electron out and repeats the measurement thousands upon thousands of times.
- (a) 4 points Each time the scientist locate the electron, will they see it as a wave or as a particle or something else?
- (b) 1 point If the scientist were also to obtain some information about the momentum of the particle, what equation relates how accurately we can determine its momentum and position?
- (c) 5 points After doing this thousands of times, the scientist finds that the probability of locating the particle in the well is:

$$P(x) = \frac{3}{8}x^2$$

What is the quantum wave function for the electron in this well, $\psi(x)$? *Just a note, this wave function isn't physically realistic.*

(d) 5 points How long is the well? *Hint: Assume that the left side of the well starts at $x = 0$ and recall that the probability over the entire well has to equal 1.*

(e) 3 points Name three effects that we see in quantum mechanics that we don't see in our everyday lives.

12. 5 points An electron moves in a straight line with constant speed $v = 1.10 \times 10^6$ m/s which has been measured with precision of 0.10%. What is the best possible precision in which its position could be measured?

13. 5 points Suppose a particle is confined to the x-axis from $x = 0$ to $x = 3$, and that its wave function is given as $\psi(x) = Ax$. What is A? *Note, this isn't a realistic function, just one that is solvable without tables of integrals.*

14. 5 points Using the function from the previous problem, find the probability of finding the particle between $x = 1$ and $x = 2$. **ALSO:** Find the average position of the particle, which is defined as

$$\int x\psi^2(x)dx$$

This is called the expectation value of the position of the particle.